

ATTRACTION AND REPULSION IN SPACETIME OF AN ELECTRICALLY CHARGED MASS-POINT

ANGELO LOINGER AND TIZIANA MARSICO

ABSTRACT. By virtue of Hilbert repulsive effect, the most external singular surface of Reißner-Weyl-Nordström metric represents an insuperable barrier for the arriving neutral particles and light rays.

PACS 04.20 – General relativity.

1. – *The metric.* – If $\gamma \equiv 1 - (2m/r) + (q^2/r^2)$, we have (see [1]÷[7]):

$$(1) \quad ds^2 = \gamma dt^2 - \gamma^{-1} dr^2 - r^2 d\omega^2 \quad ; \quad (G = c = 1) \quad ,$$

where: $d\omega^2 \equiv d\vartheta^2 + \sin^2 \vartheta d\varphi^2$; $q^2 \equiv 4\pi\epsilon^2$; and $4\pi\epsilon$ is the electric charge of the gravitating point-mass m ; $\gamma(r=0) = +\infty$; $\gamma(r=\infty) = 1$.

Two cases: $m^2 < q^2$, in particular $m = 0$; and $m^2 \geq q^2$. (For the electron we have $m^2 < q^2$).

If $m^2 < q^2$, $\gamma(r)$ is everywhere positive, with a minimal value at $r = q^2/m$: $\gamma_{\min} = 1 - (m^2/q^2)$; if $m = 0$, $\gamma_{\min} = 1$.

If $m^2 = q^2$, $\gamma(r) = [1 - (m/r)]^2$; $\gamma_{\min} = \gamma(r = m) = 0$; $\gamma_{\max} = \gamma(r = \infty) = 1$.

If $m^2 > q^2$, we can write:

$$(2) \quad r^2 \gamma(r) = \left[r - m - (m^2 - q^2)^{1/2} \right] \cdot \left[r - m + (m^2 - q^2)^{1/2} \right] \quad ;$$

let us put: $r_1 \equiv m + (m^2 - q^2)^{1/2}$ and $r_2 \equiv m - (m^2 - q^2)^{1/2}$. On the spherical surfaces $r = r_1$ and $r = r_2$, we have $\gamma(r_{1,2}) = 0$. It is easy to see that when $r_2 < r < r_1$, $\gamma(r) < 0$, but $\gamma(r < r_2) > 0$.

2. – *Radial geodesics of light-rays.* – If $ds^2 = 0 = d\omega^2$, we have $\gamma dt^2 = \gamma^{-1} dr^2$, from which:

$$(3) \quad \left(\frac{dr}{dt} \right)^2 \equiv \dot{r}^2 = \frac{(r^2 - 2mr + q^2)^2}{r^4} \quad ;$$

$[\dot{r}^2]_{r=\infty} = 1$; $[\dot{r}^2]_{r=0} = \infty$. For $m^2 > q^2$, we have $[\dot{r}^2]_{r=r_{1,2}} = 0$; and for $m^2 = q^2$, $[\dot{r}^2]_{r=m} = 0$.

The integration of [3] gives:

$$(4) \quad t + \text{const} = \pm \int \frac{r^2 dr}{r^2 - 2mr + q^2} \quad ;$$

thus, when $m^2 > q^2$:

$$(5) \quad \begin{aligned} \pm (t + \text{const}) &= r + m \ln |r^2 - 2mr + q^2| + \\ &+ \frac{2m^2 - q^2}{2(m^2 - q^2)^{1/2}} \ln \left| \frac{r - m - (m^2 - q^2)^{1/2}}{r - m + (m^2 - q^2)^{1/2}} \right| \quad ; \end{aligned}$$

this means that a light-ray which starts from an $r > r_1$, arrives at $r = r_1$ after an *infinite* time interval (and with a velocity equal to zero): we have a Hilbertian gravitational *repulsion*.

When $m^2 = q^2$:

$$(6) \quad \pm (t + \text{const}) = r - m + 2m \ln |r - m| - m^2/(r - m) \quad ;$$

for $r = m$, $\pm (t + \text{const}) = \infty$: Hilbertian repulsion, as far $m^2 > q^2$ (and a velocity $[\dot{r}]_{r=m} = 0$).

When $m^2 < q^2$, we have:

$$(7) \quad \begin{aligned} \pm (t + \text{const}) &= r + m \ln |r^2 - 2mr + q^2| + \\ &+ \frac{2m^2 - q^2}{(q^2 - m^2)^{1/2}} \cdot \arctan \left[\frac{r - m}{(q^2 - m^2)^{1/2}} \right] \quad ; \end{aligned}$$

a light-ray arrives at $r = 0$ in a finite time: gravitational attraction.

When $m = 0$:

$$(8) \quad \pm (t + \text{const}) = \int \frac{r^2 dr}{r^2 + q^2} = r - q \arctan (r/q) \quad ;$$

attraction, as for $m^2 < q^2$.

Let us compute the acceleration $\ddot{r}$:

$$(9) \quad \pm \ddot{r} = 2r^{-5} (r^2 - 2mr + q^2) (mr - q^2) \quad ;$$

of course, $[\ddot{r}]_{r=\infty} = 0$. We have:

for $m^2 > q^2$: $\ddot{r} = 0$ at $r = r_1$, repulsion; ($\ddot{r} = \infty$ at $r = 0$);
 for $m^2 = q^2$: $\ddot{r} = 0$ at $r = m$, repulsion; ($\ddot{r} = \infty$ at $r = 0$);
 for $m^2 < q^2$: $\ddot{r} = \infty$ at $r = 0$, attraction; $\ddot{r} = 0$ at $r = q^2/m$;
 for $m = 0$: $\ddot{r} = \infty$ at $r = 0$, attraction.

When $m^2 \geq q^2$ a light-ray which arrives at $r = r_1$ starting from an $r > r_1$ cannot overcome the barrier $r = r_1$: indeed, velocity $\dot{r}$ and acceleration $\ddot{r}$ are zero at $r = r_1$. (On the contrary, gravitational attraction predominates in the interval $0 \leq r < r_2$).

3. – Radial geodesics in general. – We have the following first integral:

$$(10) \quad \dot{r}^2 = \gamma^2(1 - |A|\gamma) \quad ,$$

where $A < 0$ for test-particles, and $A = 0$ for light-rays. We get from eq. (10) that

$$(11) \quad \pm \ddot{r} = \gamma r^{-2}(m - q^2 r^{-1})(2 - 3|A|\gamma) \quad .$$

The case $m^2 \geq q^2$ for $r > r_1 \equiv m + (m^2 - q^2)^{1/2}$ is interesting. When $\ddot{r} = 0$, attraction and repulsion *balance* each other. Putting $\ddot{r} = 0$ in eq. (11), we obtain

$$(12) \quad |A|\gamma = \frac{2}{3} \quad ,$$

and substituting eq. (12) in eq. (10):

$$(13) \quad \dot{r}_b^2 = \frac{1}{3}\gamma^2 \quad ;$$

a motion described by the balance velocity $\dot{r}_b$ has a zero acceleration. Where $\dot{r}^2 < \dot{r}_b^2$, there is attraction; where $\dot{r}^2 > \dot{r}_b^2$ there is *repulsion*.

At $r = r_1$, velocity $\dot{r}$ and acceleration $\ddot{r}$ are equal to zero: *no* particle, *no* light-ray can overcome this barrier. Remark that for the light-rays there is repulsion on the *whole* trajectory ($m^2 \geq q^2$).

The study of the *circular* geodesics confirms the existence of the Hilbertian repulsion: as for the Schwarzschild manifold (see [8]), there is a *minimal* value for the radial coordinate r of the test-particles, and a *unique* value for the r of light-rays ($m^2 \geq q^2$).

When $q^2 > m^2$, there is no gravitational repulsion.

A remark. If the term q^2/r^2 of $\gamma(r)$, which is originated by the mass-energy of the electrostatic field ε/r^2 , were endowed with a Newtonian nature, it would have a repulsive action. On the contrary, it has in GR an attractive effect, as it is particularly evidenced by the case $q^2 > m^2$.

4. – A coordinate choice à la Brillouin(-Schwarzschild). – For $m^2 \geq q^2$, a reasonable choice of a new radial coordinate removes the singularity at $r = 0$ of eq. (1); indeed, let us perform the following substitution:

$$(14) \quad r \rightarrow r + r_1 \quad ; \quad \left(r_1 \equiv m + (m^2 - q^2)^{1/2} \right) \quad ;$$

at $r = 0$, the new γ is zero, which is its minimal value. Our manifold is maximally extended and geodesically complete. This form of the metric is diffeomorphic to the exterior part $r \geq r_1$ of the form (1). The known generalization à la Kruskal-Szekeres of the ds^2 of eq. (1) is quite superfluous, and misrepresents the permanent gravitational field created by m and q .

For $q = 0$, substitution (14) becomes $r \rightarrow r + 2m$, *i.e.*, that substitution of the coordinate r of the *standard* metric of Schwarzschild manifold, which gives the Brillouin(-Schwarzschild) ds^2 , see [9].

In a reasonable treatment *there is no room for a real existence of black holes*. The consideration of the Brillouin(-Schwarzschild) metric reinforces the conclusion of sects. **2** and **3**.

A last remark. We think that the choice of the global time $t = x^0$ (Weyl's "kosmische Zeit" [2]) as dynamical evolution parameter for the radial geodesics of Schwarzschild, Kerr and Reißner-Weyl-Nordström manifolds is the most appropriate one from the physical point of view [8]. In particular, it allows a unique treatment of the trajectories of test-particles and light-rays, while the employment of the proper time (which is influenced by particle motion) is limited to the geodesics of the test particles.

APPENDIX

With the metric of eq. (1) the electrostatic field $F_{01}(r)$ created by the charged gravitating mass m is simply equal to ε/r^2 ("merit" of the term $r^2 d\omega^2$ in eq. (1)). As it was remarked by von Laue [6], this is an expression "die sich von der entsprechenden der Elektrostatik im Euklidischen dadurch unterscheidet, daß r nicht den natürlich gemessenen Abstand vom Ladungsträger bedeutet."

With a different form of the metric we have a more complicated expression of the e.s. field. For instance, in Bergmann's metric [7] the e.m. potential Φ is not given by $-\varepsilon/r$, but by the following formulae (where ε is equal to $-\varepsilon$ of eq. (1)):

$$(A.1) \quad \Phi = \frac{\varepsilon}{(q^2 - m^2)^{1/2}} \cdot \operatorname{arccot} \left[\frac{r - m}{(q^2 - m^2)^{1/2}} \right] ,$$

if $q^2 \equiv 4\pi\varepsilon^2 > m^2$; and by

$$(A.2) \quad \Phi = -\frac{\varepsilon}{2(m^2 - q^2)^{1/2}} \cdot \ln \left| \frac{r - m - (m^2 - q^2)^{1/2}}{r - m + (m^2 - q^2)^{1/2}} \right| ,$$

if $m^2 > q^2$. When $m^2 = q^2$, we have

$$(A.3) \quad \Phi = \frac{\varepsilon}{r - m} .$$

For $d\Phi/dr$ there is a unique formula:

$$(A.4) \quad \frac{d\Phi}{dr} = \frac{-\varepsilon}{r^2 - 2mr + q^2} .$$

It is interesting that at $r = m \pm (m^2 - q^2)^{1/2}$ the field $d\Phi/dr$ and the potential Φ have infinite values.

Bergmann's formulae make particularly evident that in GR e.m. field and gravitational field influence each other in a significant way.

REFERENCES

- [1] H. Reißner, *Ann. Physik*, **50** (1916) 106.
- [2] H. Weyl, *Ann. Physik*, **54** (1917) 117. See also *Idem*, *Raum-Zeit-Materie*, Siebente Auflage (Springer-Verlag, Berlin, *etc.*) 1988, p.257.
- [3] G. Nordström, *Versl. Akad. Wetensch. Amsterdam*, **20** (1918) 10.
- [4] C. Longo, *N. Cim.*, (VI) **15** (1918) 191.
- [5] G. B. Jeffery, *Proc. Roy. Soc.*, **99A** (1921) 123. See also A. S. Eddington, *The Mathematical Theory of Relativity*, Second Edition (Cambridge University Press, Cambridge) 1960, sect. **78**.
- [6] M. v. Laue, *Die Relativitätstheorie, Zweiter Band*, 4. Auflage (Friedr. Vieweg und Sohn, Braunschweig) 1956, sect. **47**.
- [7] P. G. Bergmann, *Introduction to the Theory of Relativity* (Prentice-Hall, Inc., Englewood Cliffs, N.Y.) 1960, p. 204 *sqq* – notice that Bergmann’s metric is different from the standard metric of our eq. (1).
- [8] D. Hilbert, *Mathem. Annalen*, **92** (1924) 1; also in *Gesammelte Abhandlungen*, Dritter Band (J. Springer, Berlin) 1935, p.258. See further A. Loinger and T. Marsico, *arXiv:0904.1578 v1* [physics.gen-ph] 9 Apr 2009 – and references [2] therein.
- [9] M. Brillouin, *Journ. Phys. Rad.*, **23** (1923) 43; for an English translation, see *arXiv:physics/0002009*, February 3rd, 2000. And references therein.

A.L. – DIPARTIMENTO DI FISICA, UNIVERSITÀ DI MILANO, VIA CELORIA, 16 - 20133 MILANO (ITALY)

T.M. – LICEO CLASSICO “G. BERCHET”, VIA DELLA COMMENDA, 26 - 20122 MILANO (ITALY)

E-mail address: `angelo.loinger@mi.infn.it`

E-mail address: `martiz64@libero.it`